Influenza (Flu)

Influenza (Flu) is a contagious respiratory illness caused by a virus. It is not the virus that mainly causes vomiting and diarrhea. The illness is self-limited with severe symptoms usually lasting 3-4 days and feeling back to normal within 2 weeks. Self-limited infections can be treated without seeing a medical provider.



SYMPTOMS: Sudden/rapid onset of symptoms

- Muscle or body aches
- Headaches
- Fever or feeling feverish with chills
- Cough

- Sore throat
- Runny or stuffy nose
- Fatigue (tiredness)
- Some people may have vomiting and diarrhea

SELF-CARE MEASURES:

- Rest
- Drink plenty of non-alcoholic fluids
- Use saline nose drops to loosen mucus
- Use Ibuprofen (Advil®) 600 mg every 6-8 hours or Acetaminophen (Tylenol®) 650 mg every 6 hours as needed to reduce fever/discomfort
- Use Oral Decongestants (Phenylephrine/Pseudoephedrine) for short-term relief of nasal congestion
- Use Nasal Decongestant (Oxymetazoline) for short-term relief of nasal congestion (do not use for more than 3 days)
- Gargle with salt water and use throat sprays/lozenges (containing Benzocaine) for throat pain
- Use heated, humidified air (if you do not have a humidifier try taking a hot shower)
- Special Note: Antiviral medication (Tamiflu®) is only recommended for patients with serious underlying medical conditions who are at risk for severe complications (such as pregnancy, immunocompromised, chronic medical conditions, or morbidly obese).

LIMIT SPREAD TO OTHERS:

- Stay home and away from others until fever-free for more than 24-hours (temperature should be less than 100 degrees Fahrenheit without medication)
- Do not go to class or eat in dining facilities
- Wash hands frequently
- Contain coughs and sneezes using the crook of your elbow
- Getting the flu vaccine is the best prevention

WHEN TO SEEK CARE FROM A MEDICAL PROVIDER:

- If you are at high risk for complications (such as pregnancy, immunocompromised, chronic medical conditions or morbidly obese).
- Fever over 102 degrees Fahrenheit that lasts for more than 3 days
- For symptoms that last over 10 days and get worse instead of better
- Shortness of breath
- Confusion or disorientation
- Severe or persistent vomiting
- Pain/pressure in your chest
- Severe sinus pain
- Very swollen glands in the neck or jaw

Loyola self-care guidelines are based on the most recent recommendations of national medical authorities.